

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/323001355>

Cassava Brown Streak Disease and Ugandan cassava brown streak virus Reported for the First Time in Zambia

Article in *Plant Disease* · February 2018

DOI: 10.1094/PDIS-11-17-1707-RE

CITATIONS

43

READS

1,027

6 authors, including:



Rabson Mulenga

Mount Makulu Central Research Station, Lusaka, Zambia

26 PUBLICATIONS 476 CITATIONS

SEE PROFILE



Laura Boykin Okalebo

BioTeam

118 PUBLICATIONS 6,471 CITATIONS

SEE PROFILE



Patrick Chiza Chikoti

Zambia Agriculture Research Institute

49 PUBLICATIONS 674 CITATIONS

SEE PROFILE



Suwilanji Sichilima

Zambia Agriculture Research Institute

7 PUBLICATIONS 289 CITATIONS

SEE PROFILE

Cassava brown streak disease and *Ugandan cassava brown streak virus* reported for the first time in Zambia

Rabson M. Mulenga¹, Laura M. Boykin², Patrick C. Chikoti¹, Suwilanji Sichilima¹, Dickson Nguni¹ and Olufemi J. Alabi³

¹Zambia Agriculture Research Institute, Mount Makulu Central Research Station, Chilanga, Lusaka, Zambia; ²The University of Western Australia, ARC Centre of Excellence in Plant Energy Biology and Molecular Sciences, Crawley, Perth 6009, Western Australia, Australia; ³Department of Plant Pathology & Microbiology, Texas A&M AgriLife Research and Extension Center, Weslaco, Texas 78596.

*Correspondence to: Olufemi J. Alabi, Department of Plant Pathology & Microbiology, Texas A&M AgriLife Research and Extension Center, Weslaco, TX 78596; email alabi@tamu.edu.

Abstract

A diagnostic survey was conducted in July 2017 in two northern districts of Zambia to investigate presence or absence of cassava brown streak disease (CBSD) and its causal viruses. Twenty-nine (29) cassava fields were surveyed and cassava leaf samples collected from 116 plants (92 symptomatic and 24 non-symptomatic). CBSD prevalence was ~79% (23/29) across fields. Mean CBSD incidence varied across fields but averaged 32.3% while mean disease severity was 2.3 on a 1-5 rating scale. RT-PCR screening of all 116 samples with one generic and two species-specific primer pairs yielded DNA bands of the expected sizes from all symptomatic plants with the generic (785 bp) and Ugandan cassava brown streak virus (UCBSV)-specific (440 bp) primers. All 24 non-symptomatic samples were negative for UCBSV and all samples tested negative with primers targeting *Cassava brown streak virus*. The complete genome of a representative isolate of UCBSV (WP282) was determined to be 9,050 nucleotide (nt) in length, minus the poly A tail. A comparative analysis of this isolate with global virus isolates revealed its nature as a sequence variant of UCBSV sharing 94/96% maximum complete polyprotein nt/amino acid identities with isolates from Malawi (MF379362) and Tanzania (FJ039520). This is the first report of CBSD and UCBSV in Zambia thus expanding the

geographical distribution of the disease and its causal virus and further reinforcing the need to strengthen national and regional phytosanitary programs in Africa.

Introduction

Cassava brown streak disease (CBSD) and cassava mosaic disease (CMD) are major constraints to cassava (*Manihot esculenta* Crantz) production in sub-Saharan Africa. Both diseases produce distinct symptoms in cassava. CMD and CBSD causal viruses are transmitted by the whitefly *Bemisia tabaci* (Gennadius) in persistent and semi-persistent manners, respectively. However, in a recent study, the DAG motif associated with aphid transmission of potyviruses was found within coat protein genes of cassava brown streak virus (CBSV) genomes and two other ipomoviruses suggestive of their potential aphid transmissibility (Ateka et al., 2017). Whereas CMD and its causal viruses have been subject of intensive studies (Thottappilly et al. 2003 and references therein; Alabi et al. 2015 and references therein), the body of knowledge on CBSD and its causal viruses is just beginning to evolve (Legg et al. 2015 and references therein).

First described in 1936 in the coastal area of Tanzania (Storey 1936), CBSD is now known to be caused by two positive-sense ssRNA viruses namely CBSV and Ugandan cassava brown streak virus (UCBSV) that belong to the genus *Ipomovirus* in the family *Potyviridae* (Mbanzibwa et al. 2009; Winter et al. 2010). The genome organization of both viruses is similar comprising a large polyprotein that encodes 10 genes but lack the helper component proteinase (HC-Pro) (Winter et al. 2010) present in most members of *Potyviridae* including in *Sweet potato mild mottle virus*, the typical *Ipomovirus* (Tugume et al. 2010). Rather, both viruses encode a unique HAM1 protein thought to be of cellular origin (Mbanzibwa et al. 2009). In cassava, CBSD symptoms due to either virus are largely indistinguishable from each other although foliar symptoms due to UCBSV are reported to be milder than those induced by CBSV (Winter et al. 2010; Patil et al. 2011). It is unclear if such phenotypic differences are due to virus- or host-dependent factors, the environment, or their interactions. However, CBSV isolates are more genetically diverse than UCBSV isolates and intraspecies, but not interspecies, recombination has been reported for both viruses (Mbanzibwa et al. 2009; Ndunguru et al. 2015; Alicai et al. 2016). Country-wide surveys have shown that UCBSV is more widespread than CBSV with confirmed reports of the former virus in eight countries compared to that of the latter in four countries (Legg et al. 2015). Additional reports exist of the presence of CBSD in several other sub-Saharan African countries

such as Angola, Gabon and Madagascar but the association of either of both viruses with these occurrences is yet to be determined (Legg et al. 2015). In addition, CBSV has been shown to have a faster rate of evolution compared to UCBSV (Alicai et al. 2016).

The landlocked country of Zambia is bordered to the north by the Democratic Republic of Congo (DRC), to the west by Angola, to the east by Malawi and Mozambique and to the south by Zimbabwe, Botswana and Namibia. So far, there are confirmed reports of UCBSV in the Democratic Republic of Congo (Mulimbi et al. 2012) and Malawi (Nichols 1950; Mbewe et al. 2015) and of CBSV in Mozambique (Hillocks et al. 2002) and Malawi (Nichols 1950; Mbewe et al. 2015). These reports heightened concerns of CBSD spread into Zambia due to the risks of inadvertent introductions via trans border movement of contaminated cassava cuttings. Hence, survey efforts were initiated in 2009 to determine the incidence and distribution of CBSD, CMD, and their causal viruses in the seven major cassava growing provinces of Zambia. These efforts led to the detection and characterization of CMD causal viruses in farmers' fields across the seven cassava growing provinces (Chikoti et al. 2013; 2015; Mulenga et al. 2016) but CBSD was reported to be absent (Chikoti et al. 2013; 2015). While areas along the long border of Zambia with Malawi were extensively surveyed, the extreme north of Luapula and Northern provinces covering Chiengwe and Kaputa districts that border the Democratic Republic of Congo were not covered. Villages and other human settlements in both districts are populated with people of mixed citizenry and it is not uncommon for cassava farmers across these largely unmanned border areas to exchange vegetative cuttings. Therefore, an intensive survey was conducted in farmers fields located in the extreme north of Luapula and Northern provinces to determine the presence or absence of CBSD and its causal viruses.

Materials and Methods

Survey and sample collection. The survey was conducted in July 2017 in the extreme northern part of Zambia covering the districts of Chiengwe in Luapula province and Kaputa in Northern province (Fig. 1). The two districts lie on the border with the Democratic Republic of Congo with no physical barrier between the two countries. Both districts were not covered in previous surveys (Chikoti et al. 2013; 2015; Mulenga et al. 2016) due to challenges associated with their accessibility. Surveys were conducted in the agriculture camps of Mwaba (4 fields), Chiengwe central (5 fields) and Chipungu (4 fields) in Chiengwe district (13 fields) and Chocha (2 fields),

Kaputa central (7 fields), Chipili (3 fields) and Kasepa (4 fields) in Kaputa district (16 fields). Cassava fields were surveyed along secondary and feeder roads encompassing areas along Lakes Mweru and Mweru Wantipa that lie at altitudes ca. 940 m above sea level (Fig. 1). Twenty-nine (29) cassava fields were assessed across both districts at 5 to 10 km intervals and each field was georeferenced using a handheld global positioning systems (GPS) device (Garmin International Inc., Olathe, KS, USA). CBSD incidence was assessed for each field as a proportion of cassava plants showing characteristic foliar and/or stem brown streak symptoms (Fig. 2) out of a total of 30 visually inspected plants along two diagonals (15 plants per diagonal) (Sseruwagi et al. 2004). Disease severity was assessed on a scale of 1-5 (Rwegasira et al. 2011). Mean incidence and severity were calculated along field diagonals for each camp and district regardless of the cultivar. To confirm root and stem diagnostic symptoms, four plants per field with foliar symptoms were uprooted, sliced and inspected for presence of root necrosis (Fig. 2G) while stems with brown streak-like symptoms (Fig. 2F) were sliced from the node upward. Symptomatic leaf samples (Figs. 2A-2E) were collected individually from four plants per field and preserved dry as per Aloyce (2013). A total of 116 (symptomatic = 92 and non-symptomatic = 24) samples were collected during the survey. All samples were packaged in labeled envelopes and transported to the Mount Makulu Central Research Station, Chilanga, Lusaka, Zambia for analysis.

RNA extraction. RNA was extracted from each dried cassava leaf sample using a previously described CTAB protocol (Alicai et al. 2016). Briefly, 50 mg of leaf tissue per sample were ground in 1 ml CTAB extraction buffer containing 2% (w/v) CTAB, 2% PVP-40, 100 mM Tris-HCl pH8.0, 25 mM EDTA, 2M NaCl, and 2% mercaptoethanol (added immediately before use). Approximately 750 μ l of the extract were pipetted into clean 2 ml microcentrifuge tubes and incubated in a water bath at 65°C for 15 min, then cooled at room temperature. Equal volume of chloroform:isoamyl alcohol (24:1 v/v) was added to the extract, tubes vortexed until content turned cloudy followed by centrifugation at 12,000 rpm for 15 min. This step was repeated twice. Finally, 600 μ l of extract were transferred into a 1.5 ml microcentrifuge tube to which one-third volume ice cold isopropanol was added followed by incubation at -20°C for 1 hr. The tubes were then centrifuged at 13000 rpm for 30 min at 4°C. The pellets were washed twice with 70% ethanol and air dried at room temperature. The dried pellet was dissolved in 50 μ l DEPC-treated

water, analyzed on a NanoDrop 2000 series spectrophotometer (ThermoFisher Scientific Inc., Waltham, MA, USA) and stored at -20°C until use.

cDNA synthesis and PCR. Two micrograms of total RNA per sample were subjected to cDNA synthesis using the Revert Aid First Strand cDNA synthesis kit (ThermoFisher Scientific Inc.) according to the manufacturer's protocol. Approximately 400 ng cDNA per sample were used in a 25 µl PCR reaction volume containing 1X DreamTaq Buffer (Life Technologies, Carlsbad, CA, USA), 0.2 mM dNTPs, 0.2 mM each sense and antisense primers, and 1 U of DreamTaq DNA Polymerase. Each of the 116 samples was screened with generic primers CPBSV1 and CPBSV2 (Aloyce 2013) and species-specific primers, CBSVF2/CBSVR7 and CBSVF2/CBSVR8 (Abarshi et al. 2012) capable of discriminating between CBSV and UCBSV, respectively. Cycling conditions were initial denaturation at 94°C for 3 min, followed by 35 cycles of 94°C for 30 s, 52°C for 30 s, 72°C for 45 s; with a final extension step of 72°C for 5 min. Additionally, each sample was screened for the presence of African cassava mosaic virus (ACMV) and East African cassava mosaic virus (EACMV), two CMD causal viruses known to be prevalent across cassava fields in Zambia (Mulenga et al. 2016; Chikoti et al. 2015). The ACMV-specific primers JSP01/02 and EACMV-specific primers EAB555F/R (Pita et al., 2001) were employed for this purpose. The PCR products were electrophoresed in 1% agarose gel pre-stained with 10 mg/ml ethidium bromide in 1X Tris-acetate-EDTA buffer and visualized using a Gel Doc XR System (Bio-Rad Laboratories, Hercules, CA, USA). An aliquot of the O'GeneRuler 1-Kb DNA Ladder (ThermoFisher Scientific, Inc.) was loaded alongside PCR amplicons for size verification.

Complete genome amplification and characterization of UCBSV. Based on results of initial screening of the 116 field collected samples, dried leaf tissues from a representative isolate (WP282) that gave DNA amplicons of the expected size with one of the primer pairs was shipped under a USDA-APHIS-PPQ permit to the Texas A&M AgriLife Research and Extension, Weslaco, Texas, USA facility for further molecular characterization. Total nucleic acid was extracted from isolate WP282 using a Spectrum Plant total RNA Kit (Sigma-Aldrich, St. Louis, MO USA) and used for cDNA synthesis with the cDNA synthesis Kit, ThermoScript™ RT-PCR System (ThermoFisher Scientific, Inc.) as per manufacturer's instructions. The genome of the CBSV causal virus present in isolate WP282 was amplified as four pieces of overlapping DNA fragments using newly designed primers (Table 1) derived

based on consensus sequence of virus isolates available in GenBank (Supplementary Table 1). The 25 µl PCR reaction volume for each primer pair consisted of 1 x PrimeSTAR GXL buffer, 0.2 mM dNTPs, primers 0.2 µM each, 1.5 U PrimeSTAR GXL DNA polymerase (Takara-Bio USA, Inc., Mountain View, CA, USA) and 2 µl of a 1:10 dilution of the template cDNA. Thermal cycling conditions were 35 cycles of 98°C for 10 s, annealing 55°C for 15 s and extension 68°C for 1 min./Kb. Correct size amplicons on a pre-stained (10 mg/ml ethidium bromide) 1% agarose gel were excised and gel eluted using Zymoclean™ Gel DNA recovery kit (Zymo Research Corp., Irvine, CA). The eluate was A-tailed and cloned into a pCR2.1 TOPO-TA vector then transformed into One Shot TOPO10 *Escherichia coli* chemically competent cells, according to manufacturer's protocol (Life Technologies, Carlsbad, CA). Two plasmids with the correct size inserts per DNA amplicon were recovered using a GenElute™ Plasmid Miniprep Kit (Sigma-Aldrich, St. Louis, MO) and initially sequenced in both directions with the M13F/R primers. Additional primers were used to walk each plasmid DNA sample (data not shown). A consensus sequence was obtained for each PCR fragment and the putative full-length viral genome was assembled using the CAP contig assembly program of the BioEdit software (Hall 1999). The genome sequence obtained from isolate WP282 was annotated based on homologies with corresponding sequences in GenBank.

Sequence analysis. The complete polyprotein and gene-specific sequences of isolate WP282 were compared with corresponding GenBank sequences (Supplementary Table 1). Sequence alignments were generated with the MUSCLE alignment program (<http://www.ebi.ac.uk/Tools/msa/muscle/>) and the alignment files employed for calculating pairwise sequence identity matrices with the BioEdit program (Hall 1999). The Simplot v.3.5.1 program (Lole et al. 1999) was used to depict a graphical representation of genome-wide sequence comparisons between isolate WP282 and selected GenBank virus isolates. The phylogenetic relationship between isolate WP282 and publicly available complete genome GenBank isolates of CBSV (14) and UCBSV (19), including infectious clones of UCBSV, was inferred using Bayesian analyses conducted with ExaBayes v.1.4.1 (Aberer et al. 2014) as described in Ndunguru et al. (2015). The polyprotein alignment file involving only UCBSV isolates was scanned for the presence of reticulate phylogenetic networks using the Neighbor-Net method as implemented in SplitsTree v4.10 (Huson and Bryant 2006). The detection of reticulate networks involving isolate WP282 rather than a tree-like topology will indicate that its

evolutionary history has been shaped by recombination (Alabi et al. 2011; Huson and Bryant 2006) as reported for CBSV causal viruses (Mbanzibwa et al. 2011; Ndunguru et al. 2015; Alicai et al. 2016). The exact nature of any potential recombination event was then investigated using the RDP4 software (Martin et al. 2015).

Results

CBSV symptoms and prevalence. Characteristic CBSV symptoms were observed in farmers' fields in the seven agriculture camps located in Chiengi (n = 3) and Kaputa (n = 4) districts. The cassava landraces commonly encountered across the 29 surveyed fields were Nshile, Kalulundyongo, Ndelekulwa, Chibungabunga and Kalilanshindo. In addition, a cassava cultivar referred to as "Research", presumably denoting that it was received from a research institute, was encountered in one field in Chiengi district. Although diverse CBSV symptoms were observed on these cultivars (Fig. 2), leaf chlorotic blotches (Figs. 2A-2B) was the most common symptom on landraces Nshile, Kalulundyongo and Ndelekulwa whereas Chibungabunga and Kalilanshindo displayed feathery veinal yellowing (Figs. 2D-2E). Of the 13 fields surveyed in Chiengi district, ~77% (10/13) had CBSV with disease incidence recorded ranging from 10% to 100%; and district-wide mean incidence of ~35% (Table 2). Similarly, ~81% (13/16) of fields surveyed in Kaputa district had CBSV with disease incidence recorded ranging from 3% to 67%; and district-wide mean incidence of 30% (Table 2). Overall CBSV prevalence across fields visited in both districts was ~79% (23/29). Moderate CBSV symptoms were observed across the 23 CBSV-affected fields regardless of the cassava cultivar with a mean severity score of 2.3 (range = 2-3.5). Characteristic CBSV root (Fig. 2G) and stem (Fig. 2F) symptoms were also observed on few plants in a limited number of locations (Table 2). In these locations, incidence of typical CBSV stem symptoms (Fig. 2F) ranged from 2.2% to 10% (low) and mean severity of root necrosis ranged from 2 to 2.3 (mild to moderate) (Table 2). Interestingly, a few plants (Chiengi = 2 and Kaputa = 3) showed symptoms of both CBSV and CMD (Fig. 2E) suggesting mixed infections of causal viruses of both diseases in these samples. Based on these results, leaf tissue samples taken from all 116 plants (symptomatic = 92; non-symptomatic = 24) were screened for presence of CBSV and CMD causal viruses.

RT-PCR analyses of CBSVs. The pair of generic CPBSV1/2 primers (Aloyce 2013) produced the expected 785 bp amplicon from all 92 symptomatic samples, while all 24 non-symptomatic

samples were negative. This indicated that each of the 92 symptomatic samples was positive for at least one of the two CBSV causal viruses. Further analysis of these 92 samples with virus-specific primers resulted in the amplification of the expected 440 bp UCBSV-specific DNA fragment from all 92 samples with the primer pair CBSVF2/CBSVR8. No DNA band was obtained from all 92 samples with the CBSV-specific primers CBSVF2/CBSVR7 indicating that UCBSV is the only CBSV causal virus present in these samples. Additionally, five (5.4%) of the 92 UCBSV-positive samples also tested positive for ACMV (4/5) and EACMV (1/5) indicating mixed infections of UCBSV with either of both cassava mosaic begomoviruses (CMBs) in these samples. Cassava plants bearing mixed infections of UCBSV and these CMBs showed more severe foliar symptoms (Fig. 2E) relative to plants with single infection of UCBSV (Figs. 2A-2D).

Determination and comparative analysis of complete genome sequence of UCBSV from Zambia. The genome of UCBSV isolate WP286 from Zambia was amplified as four pieces of overlapping DNA fragments using four pairs of newly designed primers (Table 1). Assembly of these sequence fragments resulted in a genome length of 9,050 nucleotides (nt) (GenBank Accession number MG257787) representing the complete genome of Zambian UCBSV isolate WP286, minus the poly A tail. The WP286 sequence has a 5' untranslated region (UTR) of 134 nt and a 3' UTR of 207 nt (excluding the poly A tail). The genome encodes a 2,902 amino acid (aa) long polyprotein and homology-based prediction of its proteolytic cleavage sites revealed the expected 10 individual proteins for UCBSV isolates (Fig. 3). The SimPlot analysis of an alignment file of isolate WP286 (MG257787) together with 16 GenBank isolates of UCBSV revealed that it is more closely related to isolate MLB3 (FJ039520) from Tanzania and isolate Malawi (MF379362) from Malawi relative to other GenBank isolates (Fig. 3). In pairwise comparisons, the complete polyprotein sequence of isolate WP286 shared 94% nt and 96% aa identity levels with isolates MLB3 and Malawi, its closest relatives (Table 3). In contrast, polyprotein nt/aa identity levels between isolate WP286 and the remaining GenBank isolates ranged from 86-88/92-94% (Table 3). The levels of nt/aa identities for the 10 encoded genes were similar to those of the polyprotein with a few exceptions (Table 3). For instance, whereas isolate WP286 shared similar levels of nt/aa identities with isolates MLB3 and Malawi with respect to P1, P3, CI, NIa and NIb proteins, contrasting results were obtained for the VPg, Ham1 and CP genes (Table 3). While the VPg and CP genes of isolate WP286 were more identical to

those of isolate MLB3, its Ham1 showed greater resemblance to that of isolate Malawi (Table 3). Overall, higher levels of Ham1 and CP nt/aa identities were obtained for comparisons between isolate WP282 and GenBank isolates compared to other genes indicating greater similarities between the 3' termini genes of UCBSV relative to the 5' termini genes.

Phylogenetic analysis and test for recombination. The Bayesian phylogenetic analyses confirmed the clustering of global isolates of CBSV/UCBSV into two major clades, one for each virus with isolate WP282 segregating into the UCBSV clade as expected (Fig. 4). The same five UCBSV-specific subclades identified by Alicai et al. (2016) were recovered in this study with isolate WP282 segregating into clade C along with isolates UCBSV_TZ_MLB3_(FJ039520), UCBSV_MA_43_(FN433933), and UCBSV_TZ_Tan_23_(KR108839) (Fig. 4). These results confirm the results of Alicai et al. (2016) that members of UCBSV clade C (WP282, MLB3, MA_43 and Tan_23) represent a unique UCBSV lineage. A reticulate (non-tree-like) network was detected when the UCBSV sequence alignment file was subjected to analysis with the Neighbor-Net method of the SplitsTree v4.10 program (Fig. 5). Further analysis with the different RDP4-implemented programs revealed two putative recombination events involving isolate WP282 (Table 4) besides events involving other UCBSV isolates as previously reported (Mbanzibwa et al. 2011; Ndunguru et al. 2015; Alicai et al. 2016). Results from this analysis suggest that isolate WP282 arose potentially as a consequence of genetic recombination.

Discussion

CBSD and CMD are no doubt the most serious constraints to sustainable production of cassava in sub-Saharan African countries. Unlike CMD which is endemic in all sub-Saharan African cassava-producing countries, the geographical distribution of CBSD and its causal viruses have so far been limited largely to the southeastern axis of the African continent. At present, there are confirmed reports of CBSD occurrence in nine countries including Uganda, Kenya, Malawi, Tanzania, Mozambique, Rwanda, Burundi, Democratic Republic of Congo and the Indian Oceanic island of Mayotte (Legg et al. 2015 and references therein). Given the proximity of Zambia to some of these countries including the fact that it shares extensive land borders with Malawi and the Democratic Republic of Congo (Fig. 1) it seemed inevitable that CBSD would eventually spread into Zambia. Consequently, CBSD and its causal viruses are considered priorities in surveillance efforts by scientists from the Zambian Agricultural Research Institute

and the International Institute of Tropical Agriculture. Interestingly, despite extensive surveys (Chikoti et al. 2013; 2015), CBSD and its causal viruses have not been found in Zambia prior to this study thus prompting the current survey effort. With the mean incidences of ~32% across ~79% (23/29) farmer's fields in the surveyed districts, it is evident that CBSD is relatively widespread in both Chiengi (Luapula province) and Kaputa (Northern province) districts of Zambia. In these fields, CBSD was documented on commonly grown cassava landraces and on the cultivar that farmers referred to as "research". Due to the annual growth pattern of cassava, it is difficult to estimate how long CBSD had been present but a ~79% (23/29) prevalence and ~32% mean disease incidence across surveyed fields and all six encountered cassava cultivars would suggest that local spread has occurred over time albeit at low rates. The only logical explanation for CBSD incursion into Zambia is that the disease was probably initially introduced inadvertently via farmer-to-farmer exchange of contaminated cassava planting material rather than by the whitefly vector. Subsequently, the combination of low whitefly population in most of the cassava growing provinces in Zambia (Chikoti et al. 2015) and low *B. tabaci* transmission efficiency of CBSVs (Maruthi et al. 2005) contributed to a limited CBSD spread encountered during the survey. This is in addition to the universally accepted view that long distance spread of whitefly transmitted viruses is caused by human movement of infected planting materials from regions where disease occurs to new areas (Legg 1999; Mbewe et al. 2015).

The detection of UCBSV alone from all 92 symptomatic plants sampled in this study support the view that it is more widespread than CBSV (Bigirimana et al. 2011; Mbanzibwa et al. 2011; Mulimbi et al. 2012; Mbewe et al. 2015; Michel et al. 2016). The exact reasons for the widespread nature of UCBSV relative to CBSV remains unclear especially against the background that isolates of the latter virus are more virulent than those of the former (Ogwok et al. 2015; Alicai et al. 2016). While initial reports point to an altitude dependent prevalence of CBSV and UCBSV (Alicai et al. 2007; Mbanzibwa et al. 2009), an altitude independent occurrence of both viruses has also been reported recently (Ndunguru et al. 2015; Alicai et al. 2016). Hence, studies of the comparative epidemiology of both viruses and the relative ability of the whitefly vector to acquire and transmit them are warranted in order to better understand factors responsible for the widespread nature of UCBSV than CBSV.

The majority (87/92) of the symptomatic UCBSV-positive cassava plants were singly infected with the virus and exhibited mild symptoms phenotype (Figs. 2A-2D) in agreement with

previous reports (Mbanzibwa et al. 2009; Legg et al. 2015; Ogwok et al. 2015). These results indicated prevalence of single infection of UCBSV in the surveyed fields. However, five symptomatic plants of the landraces Chibungabunga and Kalilanshindo from three cassava fields located in Kaputa district showed severe symptoms (Fig. 2E). Further evaluation of these plants showed mixed infections of UCBSV with ACMV or EACMV thus suggestive of synergism between the co-infecting viruses. Root and stem symptoms (Figs. 2F-2G) were also observed in some of the disease-affected plants in the two districts, Chiengwe and Kaputa consistent with symptoms documented in previous studies (Nichols 1950; Mbewe et al. 2014; Patil and Fauquet 2014) and indicating that these phenotypes directly correlated to negative impacts of CBSD are also present under Zambian growing conditions. However, foliar symptoms predominated in all 23 CBSD-affected fields in agreement with previous reports which indicated that foliar CBSD symptoms are the most common and streaks on stems and root necrosis are irregular in cassava plants naturally infected with CBSD causal viruses (Ogwok et al. 2010; Mbewe et al. 2014; Ndunguru et al. 2015; Alicai et al. 2016).

The genome characterization of isolate WP286 from Zambia and its clustering pattern (Fig. 4) further support the distinct nature of the UCSBV clade C isolates identified in a recent study (Ndunguru et al. 2015; Alicai et al. 2016). In this clade, isolate WP286 showed contrasting patterns of gene-specific sequence similarities and differences with other members indicating its distinctness as a sequence variant of UCBSV. Interestingly, the observed contrasting patterns of sequence similarities between WP282 and its closest relatives occurred in the 3' distal *Ham1* and *CP* genes whereas similar levels of sequence similarities were observed for other regions of the genome (Table 3; Fig. 3). These observations point to potential genetic exchanges occurring among natural populations of UCBSV perhaps as a consequence of an expansion in its geographical distribution (Mbandizwa et al. 2011). Notably the sequence of recently generated UCBSV infectious clones Kikombe (KX753356) and Kikombe-IC (KX753357) (Nanyiti, Unpublished) differ considerably from those of clade C isolates (Table 3; Figs. 3 and 4) hence there may be need to generate additional clade-specific infectious clones for use by breeders and scientists in breeding and reverse genetics studies.

A pertinent question of interest to national and regional regulatory agencies is regarding the possible route of spread of CBSD into Zambia. The relatively high level of similarity of isolate WP282 with UCBSV isolates from Malawi and Tanzania (Table 3, Figs. 3 and 4) suggests

possible incursion of the disease from either country into Zambia. However, recent intensive and extensive surveys conducted especially in farmer's fields along the long border of Zambia with Malawi reported absence of CBSD in these areas (Chikoti et al. 2013; 2015). Therefore, it is logical to propose that isolate WP282 was probably introduced into Zambia from Tanzania. The support for this inference is found in higher levels of sequence similarities obtained between isolate WP282 and Tanzanian isolate MLB3 in nine of ten genes compared to eight of ten genes with isolate Malawi (Table 3). Interestingly, isolate MLB3 was sourced from the Kagera region of Northwestern Tanzania which is bordered to the west by Lake Tanganyika, a freshwater lake that supports commerce and human movement across countries Tanzania, Democratic Republic of the Congo, Burundi, and Zambia. It is also conceivable that UCBSV introduction into Zambia came from the Democratic Republic of Congo with whom the surveyed districts share land border (Fig. 1). However, no genomic data is available from the Democratic Republic of Congo hence future studies are needed to help us track the true origin of UCBSV introduction into Zambia. Regardless, there exists very high risk of CBSD spread post-introduction to other cassava-growing provinces of Zambia via movement of infected stems and/or whiteflies if urgent exclusionary measures are not taken by relevant regulatory agencies in the country. Therefore, our results point to the need to strengthen phytosanitary programs and intensify farmer extension and outreach efforts in the region to stem the tide of disease spread via movement of plant germplasm.

In conclusion, we have detected the occurrence of CBSD for the first time in Zambia and the association of a sequence variant of UCBSV with the disease in affected fields. Efforts have been initiated by the Zambian Agriculture Research Institute (ZARI) to destroy cassava plants in all affected fields to avert further spread inland to cassava fields in the seven major cassava producing provinces of the country.

Acknowledgement

This project was supported with funds from the Agricultural Productivity Program for Southern Africa (APPSA) Zambia, awarded to Zambia Agriculture Research Institute (ZARI), Grant number P094183-IDA52050 (Subproject: CV-P01-2016). Travel funds for Mr. Rabson Mulenga's visit to the Texas A&M AgriLife Research and Extension Center, Weslaco facility was provided by the Norman E. Borlaug Leadership Enhancement in Agriculture Program

(Borlaug-LEAP) funded by USAID. Computational resources provided by the Pawsey Supercomputing Centre with funding from the Australian Government and the Government of Western Australia supported this work. We are also grateful to the management of the Texas A&M AgriLife Research and Extension Center, Weslaco for facility support and to Ms. Aisha Zongoma (Ahmadu Bello University, Zaria, Nigeria) for technical assistance.

References

- Abarshi, M. M., Mohammed, I. U., Jeremiah, S. C., Legg, J. P., Lava Kumar, P., Hillocks, R. J., and Maruthi, M. N. 2012. Multiplex RT-PCR assays for the simultaneous detection of both RNA and DNA viruses infecting cassava and the common occurrence of mixed infections by two cassava brown streak viruses in East Africa. *J. Viro. Methods*. 179:176-184.
- Aberer, A. J., K. Kobert, and A. Stamatakis. 2014. ExaBayes: Massively Parallel Bayesian Tree Inference for the Whole-Genome Era. *Molecular Biology and Evolution* 31:2553-2556.
- Alabi, O. J., Al Rwahnih, M., Karthikeyan, G., Poojari, S., Fuchs, M., Rowhani, A., and Naidu, R. A. 2011. *Grapevine leafroll-associated virus 1* occurs as genetically diverse populations. *Phytopathol.* 101:1446-1456
- Alabi, O. J., Mulenga, R. M., and Legg, J. P. 2015. Cassava mosaic. In: Tennant P, Fermin G, eds. CABI International: Virus diseases of tropical and subtropical crops. CABI UK, 56-72.
- Alicai, T., Ndunguru, J., Sseruwagi, P., Tairo, F., Okao-Okuja, G., Nanvubya, R., Kiiza, L., Kubatko, L., Kehoe, M., and Boykin, M. L. 2016. Cassava brown streak virus has a rapidly evolving genome: implications for virus speciation, variability, diagnosis and host resistance. *Sci. Rep.* 6:36164 DOI: 10.1038/srep36164.
- Alicai, T., Omongo, C. A., Maruthi, M. N., Hillocks, R. J., Baguma, Y., Kawuki, R., Bua, A., Otim-Nape, G.W., and Colvin, J. 2007. Re-emergence of cassava brown streak disease in Uganda. *Plant Dis.* 91:24-29.
- Aloyce, 2013. Development and evaluation of efficient diagnostic tools for cassava mosaic and cassava brown streak diseases. MSc. Thesis, The University of Witwatersrand, Johannesburg Republic of South Africa.
- Ateka, E., Alicai, T., Ndunguru, J., Tairo, F., Sseruwagi, P., Kiarie, S., Makori, T., Kehoe, M. A., and Boykin, L. M. 2017. Unusual occurrence of a DAG motif in the Ipomovirus *Cassava brown streak virus* and implications for its vector transmission. *PLoS ONE* 12(11):e0187883.
- Bigirimana, S., Barumbanze, P., Ndayihanzamaso, P., Shirima, R., and Legg, J. P. 2011. First report of cassava brown streak disease and associated *Ugandan cassava brown streak virus* in Burundi. *New Dis. Rep.* 24:26.

- Chikoti, P. C., Ndunguru, J., Melis, R., Tairo, F., Shanahan, P., and Sseruwagi, P. 2013. Cassava mosaic disease and associated viruses in Zambia: occurrence and distribution. *Int. J. Pest Mgt.* 59:63-72.
- Chikoti, P.C., Tembo, M., Chisola, M., Ntawuruhunga P., and Ndunguru, J. 2015. Status of cassava mosaic disease and whitefly population in Zambia. *Afr. J. Biotech.* 14:2539-2546.
- Hall, T. A. 1999. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids. Symp. Ser.* 41:95-98.
- Hillocks, R. J., Thresh, J. M., Tomas, J., Botao, M., Macia, R., Xavier, R. 2002. Cassava brown streak disease in northern Mozambique. *Int. J. Pest Manage.* 48:179-182.
- Huson, D. H., and Bryant, D. 2006. Application of phylogenetic networks in evolutionary studies. *Mol. Biol. Evol.* 23:254-267.
- Legg, J. P. 1999. Emergence, spread and strategies for controlling the pandemic of cassava mosaic virus disease in east and central Africa. *Crop Prot.* 18:627-637.
- Legg, J. P., Lava Kumar., Makesh Kumar, T., Tripathi, L., Ferguson, M., Kanju, E., Ntawuruhunga, P., and Cuellar, W. 2015. Cassava Virus Diseases: Biology, Epidemiology, and Management. *Adv. In Virus Res.* 91:85-142.
- Lole, K. S., Bollinger, R. C., Paranjape, R. S., Gadkari, D., Kulkarni, S. S., Novak, N. G., Ingersoll, R., Sheppard, H. W., and Ray, S. C. 1999. Full-length human immunodeficiency virus type 1 genomes from subtype C-infected seroconverters in India, with evidence of intersubtype recombination. *J. Virol.* 73(1):152-60.
- Martin, D. P., Murrell, B., Golden, M., Khoosal, A., Muhire, B. 2015. RDP4: detection and analysis of recombination patterns in virus genomes. *Virus Evol.* 1(1):1-5.
- Maruthi, M. N., Hillocks, R. J., Mtunda, K., Raya, M. D., Muhanna, M., Kiozia, H., Rekha, A. R., Colvin, J., Thresh, J. M., 2005. Transmission of *Cassava brown streak virus* by *Bemisia tabaci* (Gennadius). *J. Phytopathol.* 153:307-312.
- Mbanzibwa, D. R., Tian, Y. P., Tugume, A. K., Mukasa, S. B., Tairo, F., Kyamanywa, S., Kullaya, A., and Valkonen, J. P. T. 2009. Genetically distinct strains of *Cassava brown streak virus* in the Lake Victoria basin and the Indian Ocean coastal area of East Africa. *Arch. Virol.* 154:353-359.
- Mbanzibwa, D. R., Tian, Y. P., Tugume, A. K., Patil, B. L., Yadav, J. S., Bagewadi, B., Abarshi, M. M., Alicai, T., Changade, W., Mkumbira, J., Muli, M. B., Mukasa, S. B., Tairo, F., Baguma, Y., Kyamanywa, S., Kullaya, A., Maruthi, M. N., Fauquet, C. M., and Valkonen, J. P. T. 2011. Evolution of cassava brown streak disease-associated viruses. *J. Gen. Virol.* 92(4):974-87.
- Mbewe, W., Kumar, P. L., Changadeya, W., Ntawuruhunga, P., and Legg, J. 2015. Diversity, distribution and effects on cassava cultivars of cassava brown streak viruses in Malawi. *J. Phytopathol.* 163(6):433-43.

- 445 Michel, R-C., David, T, Thomas C., Charles, J., Delphine M., Katia, J., Liachourutu, A. K. A.,
 446 Bruno H., Bernard R., Legg J. P., and Lett, J-M. 2016. Occurrence of cassava brown
 447 streak disease and associated *Ugandan cassava brown streak virus* in Mayotte Island
 448 (Mozambique Channel). In: Electronic proceedings of the World Congress on Root and
 449 tuber crops. Naming: WCRTC, China. <http://www.Gcp21.org/wcrtc/504>.
- 450 Mulenga, R. M., Legg, J. P., Ndunguru, J., Chikoti, P. C., Miano, D. W., Mutitu, W. E., and
 451 Alabi, O. J. 2016. Survey, molecular detection, and characterization of geminiviruses
 452 associated with cassava mosaic disease in Zambia. *Plant Dis.* 100:1379-1387.
- 453 Mulimbi, W., Phemba, X., Assumani, B., Kasereka, P., Muyisa, S., Ugentho, H., Reeder, R.,
 454 Legg, J. P., Laurenson, L., Weekes, R., and Thom, F. E. F. 2012. First report of *Ugandan*
 455 *cassava brown streak virus* on cassava in Democratic Republic of Congo. *New Dis. Rep.*
 456 26:11.
- 457 Ndunguru, J., Sseruwagi, P., Tairo, F., Stomeo, F., Maina, S., Djinkeng, A., Kehoe, M., and
 458 Boykin, M. L. 2015. Analyses of twelve new whole genome sequences of cassava brown
 459 streak viruses and Ugandan cassava brown streak viruses from East Africa: diversity,
 460 supercomputing and evidence for further speciation. *PLoS ONE*10(10): e0139321.
- 461 Nichols, R. F. J., 1950. The brown streak disease of cassava: distribution, climatic effects and
 462 diagnostic symptoms. *East Afr. Agric. J.* 15:154-160.
- 463 Ogowok, E., Alicai, T., Rey, M. E. C., Beyene, G., and Taylor, N. J. 2015. Distribution and
 464 accumulation of cassava brown streak viruses within infected cassava (*Manhot esculenta*)
 465 plants. *Plant Pathol.* 64(5):1235-1246.
- 466 Ogowok, E., Patil, B. L., Alicai, T., and Fauquet, C. M. 2010. Transmission studies with Cassava
 467 brown streak Uganda virus (*Potyviridae: Ipomovirus*) and its interaction with abiotic and
 468 biotic factors in *Nicotiana benthamiana*. *J. Virol. Methods.* 169:296-304.
- 469 Patil, B. L., and Fauquet, C. M. 2014. Light intensity and temperature affect systemic spread of
 470 silencing signal in transient agroinfiltration studies. *Mol. Plant Pathol.* 16(5):484-94.
- 471 Patil, BL., Legg, J.P., Kanju, E., Fauquet, C.M. 2015. Cassava brown streak disease: a threat to
 472 food security in Africa. *J. Gen. Virol.* 96:956-968.
- 473 Patil, B. L., Ogowok, E., Wagaba, H., Mohammed, I. U., Yadav, J. S., Bagewadi, B., Taylor, N.
 474 J., Kreuze, J. F., Maruthi, M. N., Alicai, T., and Fauquet, C. M., 2011. RNAi-mediated
 475 resistance to diverse isolates belonging to two virus species involved in cassava brown
 476 streak disease. *Mol. Plant Pathol.* 12:31-41.
- 477 Rwegasira, G. M., Momanyi, G., Rey, M. E. C., Kahwa, G., and Legg, J. P. 2011. Widespread
 478 occurrence and diversity of *Cassava brown streak virus* (*Potyviridae: Ipomovirus*) in
 479 Tanzania. *Phytopathol.* 101:1159-1167.
- 480 Sseruwagi, P., Sserubombwe, W. S., Legg, J. P., Ndunguru, J., Thresh, J. M. 2004. Methods of
 481 surveying the incidence and severity of cassava mosaic disease and whitefly vector
 482 populations on cassava in Africa: a review. *Virus Res.* 100:129-142.

- 483 Storey, H. H., 1936. Virus diseases of East African plants. VI-A progress report on studies of the
484 disease of cassava. East Afr. Agric. J. 2:34-39.
- 485 Thottappilly, G., Thresh, J.M., Calvert, L.A and Winter, S. 2003. Cassava. In “Virus and virus-
486 like diseases of major crops in developing countries”. Pp 107-165, Kluwer Academic
487 Publishers.
- 488 Tugume, A. K., Cuéllar, W. J., Mukasa, S. B., and Valkonen, J. P. T. 2010. Molecular genetic
489 analysis of virus isolates from wild and cultivated plants demonstrates that East Africa is a
490 hotspot for the evolution and diversification of *Sweet potato feathery mottle virus*. Mol.
491 Ecol. 19:3139-3156.
- 492 Winter, S., Koerbler, M., Stein, B., Pietruszka, A., Paape, M., and Butgereitt, A., 2010. Analysis
493 of cassava brown streak viruses reveals the presence of distinct virus species causing
494 cassava brown streak disease in East Africa. J. Gen. Virol. 91:1365-1372.

Table legends.

Table 1. Oligonucleotides used for amplification of complete genome of Ugandan cassava brown streak virus (UCBSV) isolate WP282 from Zambia.

Table 2. Incidences of foliar, stem and root symptoms of cassava brown streak disease (CBSD) recorded in Zambia based on a diagnostic survey conducted in July 2017 in two northern districts of Luapula and Northern provinces of the country.

Table 3. Percent nucleotide (nt)/amino acid (aa) sequence identities between polyprotein and encoded genes of Zambian isolate WP282 of *Ugandan cassava brown streak virus* and global isolates of the virus.

Table 4. Putative recombination events involving *Ugandan cassava brown streak virus* isolate WP282.

Supplementary Table 1. Sequences assembled from the NCBI database and used to design primers UCBSVv502/ UCBSVc3182 and UCBSVv6456/ UCBSVc9055 (Table 1).

Figure legends.

Figure 1. A partial map of Africa showing Zambia and adjoining countries and survey locations targeted in this study; Chiengwe and Kaputa districts of Luapula and Northern provinces of Zambia. Red and black dots denote cassava fields with and without cassava brown streak disease (CBSD) symptoms, respectively. LP = Luapula province, LSP = Lusaka province, EP = Eastern province, NP = Northern province, CBP = Copperbelt province, CP = Central province, WP = Western province, SP = Southern province, and NWP = Northwestern province.

Figure 2. Symptoms of cassava brown streak disease (CBSD) observed on naturally infected cassava plants in Chiengwe (Luapula province) and Kaputa (Northern province) districts of Zambia. The observed foliar symptoms include chlorotic blotches (A & B), mottling (C), and feathery vein chlorosis (D & E). Severely affected plants may also show brown streaks on stems (F) and root necrosis (G). In addition to CBSD symptoms, few plants also displayed symptoms of mosaic and leaf distortion typical of cassava mosaic disease (Fig. 2E).

Figure 3. Sliding-window Simplot graph showing genome-wide comparisons of isolate WP282 of Ugandan cassava brown streak virus (UCBSV) with other isolates of the virus. The Simplot graph was generated using multiple sequence alignments of UCBSV isolates with a window size

of 200 nt and a step size of 20 nts (Kimura-2 parameter). A diagrammatic representation of the UCBSV genome with locations of the encoded proteins from 5' to 3' is shown above the graph. Details of genome organizations of UCBSV can be found in Mbanzibwa et al. (2011).

Figure 4. Bayesian analysis depicting evolutionary relationships between Ugandan cassava brown streak virus (UCBSV) isolate WP282 from Zambia and global isolates of the virus and Cassava brown streak virus (CBSV). The analysis was conducted using ExaBayes version 1.4.1 (Aberer et al., 2014) essentially as described in Ndunguru et al. (2015).

Figure 5. SplitsTree reticulate (non-tree-like) networks analyses of complete polyprotein sequences of *Ugandan cassava brown streak virus*. The network analyses were performed using the Neighbor-Net method as implemented in SplitsTree v4.10 (Huson and Bryant, 2006). GenBank accession numbers of the isolates are provided on Table 3.

Table 1. Oligonucleotides used for amplification of complete genome of Ugandan cassava brown streak virus (UCBSV) isolate WP282 from Zambia (MG257787).

Name ^a	Sequence (5'-3') ^b	Size (bp)
UCBSVv1	AAAAATCACATACTCATgACATAA	549
UCBSVc549	gCTgTAgCCTTCgTTCTAC	
UCBSVv502	AgTYgCTTgTgCCTTTg	2681
UCBSVc3182	CAAgggTAACACCATTTC	
UCBSVv2945	ggAACATCTgTgTTTggTgATg	3619
UCBSVc6563	CTTCACTTTCCATTACAgCCTTg	
UCBSVv6456	CTAAGaAGCACCGTggAAAT	2600
UCBSVc9055	CTACACCRAACARAaggATATg	

^aSmall case letters v and c indicate virion and complementary sense primers, respectively. Numerals following both symbols indicate genome position of the starting nucleotide for each primer based on isolate WP282.

^bY = C+T; R = A+G; D = G+A+T; M = A+C; W = A+T. Primers UCBSVv502/ UCBSVc3182 and UCBSVv6456/ UCBSVc9055 were designed based on consensus sequence of alignments derived from 17 GenBank isolates (Supplementary Table 1) while the remaining two primer pairs are specific to isolate WP282.

Table 2. Incidences of foliar, stem and root symptoms of cassava brown streak disease (CBSD) recorded in Zambia based on a diagnostic survey conducted in July 2017 in the northern districts of Chiengge (Luapula province) and Kaputa (Northern province) of the country.

District	Camp	Mean altitude (masl)	No. of fields (CBSD+ve/CBSD-ve) ^a	Mean CBSD incidence ^b		Mean root CBSD ^c	
				Foliar (%)	Stem (%)	Root necrosis	Severity
Chiengge	Chiengge central	1066	5 (2/3)	35	0	0	NS
	Mwabu	947	4 (4/0)	25	4.2	0	NS
	Chipungu	947	4 (4/0)	75	10	0.19	2.15
Kaputa	Kaputa central	975	7 (6/1)	35.	0	0.14	2.15
	Chocha	974	2 (1/1)	33.5	0	0	NS
	Chipili	955	3 (3/0)	40	2.2	0	NS
	Kasepa	978	4 (3/1)	25	0	0	NS

^aCBSD+ve = presence of cassava plants showing CBSD symptoms; CBSD-ve = absence of cassava plants showing CBSD symptoms

^b% incidence = (number of symptomatic plants/total number of plants evaluated) x 100

^cMean root necrosis was calculated using the formula (x/n), where x = number of roots with necrosis, n = total number of tubers inspected. Only four plants were sampled per field and number of root tubers per plant ranged from 2 to 6. NS = No CBSD root necrosis symptoms. Mean root CBSD severity range was calculated as previously described by Rwegasira et al. (2011).

Table 3. Percent nucleotide (nt)/amino acid (aa) sequence identities between polyprotein and encoded genes of Zambian isolate WP282 of *Ugandan cassava brown streak virus* (MG257787) and global isolates of the virus.

Isolate	Accession no.	Polyprotein	P1	P3	6K1	CI	6K2	VPg	NIa	NIb	Ham1	CP
MLB3	FJ039520	94/96	94/94	95/96	96/96	95/99	97/98	95/99	95/97	94/97	92/92	94/95
Malawi	MF379362	94/96	94/94	95/96	94/100	95/99	97/98	94/96	95/97	95/97	94/94	91/94
F25S6.S10N	KR911725	88/93	92/93	85/87	81/96	88/98	83/94	84/91	85/95	86/93	90/89	91/94
Kikombe	KX753356	86/92	83/88	84/86	85/96	86/97	85/96	85/90	84/92	86/93	90/89	91/94
Kikombe-IC	KX753357	86/92	83/88	85/86	85/96	86/97	85/96	85/90	84/92	86/93	90/89	91/94
F10S2.S20C	KR911722	86/93	83/88	84/86	85/96	88/98	82/96	85/92	85/96	85/93	89/88	91/96
Ma42	FN433932	86/93	84/90	85/86	79/92	88/98	82/92	85/90	85/96	86/94	90/91	91/93
Ma43	FN433933	87/93	84/90	85/86	80/94	87/98	82/92	85/90	85/96	86/95	90/92	91/95
Ke125	FN433930	87/93	83/88	86/88	84/96	88/97	81/96	84/90	85/94	86/94	90/91	92/95
Ug23	FN434109	87/94	83/89	85/87	81/96	87/98	84/96	85/91	85/95	87/95	92/91	92/96
Uganda:Nam	HM181930	87/93	84/89	84/86	83/96	88/97	83/94	83/90	85/94	86/93	89/90	91/95
Ugandan	FJ185044	87/96	84/89	84/86	83/96	88/97	83/94	83/90	85/94	86/93	89/90	91/95
F16S4.S7W	KR911723	87/94	84/90	85/87	83/98	88/98	85/96	84/91	85/94	86/95	90/91	90/95
F17S3.S2W	KR911724	87/93	84/90	85/87	83/96	88/98	85/96	85/91	85/94	86/95	90/91	91/95
UG:Kab	HG965222	87/93	84/90	84/86	83/96	88/98	85/96	85/90	86/95	85/94	89/89	91/95
Ke54	FN433931	87/93	83/89	84/87	81/96	88/97	85/96	85/91	85/95	86/95	91/91	90/93

Descriptions of Ugandan cassava brown streak virus (UCBSV) encoded proteins can be found in Mbanzibwa et al. (2011).

Table 4. Putative recombination events involving *Ugandan cassava brown streak virus* isolate WP282 (MG257787).

Putative recombinant isolate	GenBank accession no. ^a	'Parent-like' isolates ^b		Method ^c	P-value ^d
		Major	Minor		
WP282	6919 (NIb), 7772 (CP)	Unknown	Ma43 (90%)	M, C , 3S	7.514 x 10 ⁻⁰⁴
Malawi	7796 (CP), 31 (P1)	WP282 (95%)	Ma43 (98%)	R, G, B , M, C, S, 3S	5.645 x 10 ⁻³⁵

^aGenomic region corresponding to the putative recombination breakpoints are shown in parenthesis.

^bThe percent similarity value shared by the putative recombinant with each parent-like isolate is shown in parenthesis.

^cThe different recombination detection used in the RDP3 analyses are R, RDP; G, GENECONV; B, BOOTSCAN; M, MAXCHI; C, CHIMAERA; S, SISCAN; and 3S, 3SEQ. Only events supported by at least three of the seven RDP4-implemented methods (Martin et al. 2005) are reported.

^dThe support probability (*P*-value) for the method in bold font is shown.




Figure 1. A partial map of Africa showing Zambia and adjoining countries and survey locations targeted in this study; Chiengwe and Kaputa districts of Luapula and Northern provinces of Zambia. Red and black dots denote cassava fields with and without cassava brown streak disease (CBSD) symptoms, respectively. LP = Luapula province, LSP = Lusaka province, EP = Eastern province, NP = Northern province, CBP = Copperbelt province, CP = Central province, WP = Western province, SP = Southern province, and NWP = Northwestern province.

992x597mm (96 x 96 DPI)




Figure 2. Symptoms of cassava brown streak disease (CBSD) observed on naturally infected cassava plants in Chiengwe (Luapula province) and Kaputa (Northern province) districts of Zambia. The observed foliar symptoms include chlorotic blotches (A & B), mottling (C), and feathery vein chlorosis (D & E). Severely affected plants may also show brown streaks on stems (F) and root necrosis (G). In addition to CBSD symptoms, few plants also displayed symptoms of mosaic and leaf distortion typical of cassava mosaic disease (E).

197x125mm (150 x 150 DPI)





Figure 3. Sliding-window Simplot graph showing genome-wide comparisons of isolate WP282 of Ugandan cassava brown streak virus (UCBSV) with other isolates of the virus. The Simplot graph was generated using multiple sequence alignments of UCBSV isolates with a window size of 200 nt and a step size of 20 nts (Kimura-2 parameter). A diagrammatic representation of the UCBSV genome with locations of the encoded proteins from 5' to 3' is shown above the graph. Details of genome organizations of UCBSV can be found in Mbanzibwa et al. (2011).

228x117mm (150 x 150 DPI)






Figure 5. SplitsTree reticulate (non-tree-like) networks analyses of complete polyprotein sequences of *Ugandan cassava brown streak virus*. The network analyses were performed using the Neighbor-Net method as implemented in SplitsTree v4.10 (Huson and Bryant, 2006). GenBank accession numbers of the isolates are provided on Table 3.

199x121mm (131 x 131 DPI)

Supplementary Table 1. Sequences assembled from the NCBI database and used to design primers UCBSVv502/ UCBSVc3182 and UCBSVv6456/ UCBSVc9055 (Table 1).

Virus	GenBank accession no.	Reference
UCBSV	HM181930	Monger et al., 2010
UCBSV	FN433930	Winter et al., 2010
UCBSV	FN433931	Winter et al., 2010
UCBSV	FN433932	Winter et al., 2010
UCBSV	FN434109	Winter et al., 2010
UCBSV	KR108835	Ndunguru et al., 2015
UCBSV	KR108836	Ndunguru et al., 2015
UCBSV	KR108837	Ndunguru et al., 2015
UCBSV	KR108838	Ndunguru et al., 2015
UCBSV	KR108839	Ndunguru et al., 2015
UCBSV	KX753357	Nanyiti et al., 2016
UCBSV	KX753356	Nanyiti et al., 2016
UCBSV	KR911727	Kathurima et al., 2016
UCBSV	FN433933	Winter et al., 2010
UCBSV	KR911728	Kathurima et al., 2016
UCBSV	KR911726	Kathurima et al., 2016